

Contents lists available at [SciVerse ScienceDirect](http://www.sciencedirect.com)

## Journal of Insect Physiology

journal homepage: [www.elsevier.com/locate/jinsphys](http://www.elsevier.com/locate/jinsphys)Differences in microRNAs and their expressions between foraging and dancing honey bees, *Apis mellifera* L.Li Li<sup>a</sup>, Fang Liu<sup>a</sup>, Wenfeng Li<sup>a</sup>, Zhiguo Li<sup>a</sup>, Jiao Pan<sup>a</sup>, Limin Yan<sup>a</sup>, Shaowu Zhang<sup>a,b</sup>, Zachary Y. Huang<sup>c,\*</sup>, Songkun Su<sup>a,d,\*</sup><sup>a</sup> College of Animal Sciences, Zhejiang University, Hangzhou, China<sup>b</sup> ARC Centre of Excellence in Vision Science, Research School of Biology, College of Medicine, Biology and Environment, The Australian National University, Canberra, Australia<sup>c</sup> Department of Entomology, Michigan State University, East Lansing, MI 48824, USA<sup>d</sup> College of Bee Science, Fujian Agriculture and Forestry University, Fuzhou, China

## ARTICLE INFO

## Article history:

Received 2 May 2012

Received in revised form 15 August 2012

Accepted 16 August 2012

Available online 21 September 2012

## Keywords:

*Apis mellifera ligustica*

Foraging behavior

Dancing behavior

MicroRNAs

## ABSTRACT

Many studies have established that microRNAs (miRNAs) regulate gene expression in various biological processes in mammals and insects including honey bees. Dancing behavior is a form of communication unique to honey bees. However, it remains unclear which miRNAs regulate the dancing behavior in honey bees, and how. In the present study, total small RNAs (sRNAs) in *Apis mellifera* foragers and dancers were extracted and analyzed by a Solexa Sequencer to determine differentially expressed miRNAs. A small percentage (12.62%) of the unique sRNAs (the number of sequence types) were shared between foragers and dancers, but their expression accounted for 92.92% of the total sRNAs (the number of all sequence reads), and the length of them centered around 22 nt. Out of 58 previously identified miRNAs, 54 were present in both foragers and dancers and most of them were down-regulated in dancers. The fold-changes of ame-miR-34, ame-miR-210, ame-miR-278 and ame-miR-282 were higher than 2. 86 and 104 novel miRNAs were detected in foragers and dancers, respectively. Furthermore, two known miRNAs (ame-miR-278 and ame-miR-282) were confirmed, by qPCR, to have lower expressions in dancers. The target genes of ame-miR-278 and ame-miR-282 were associated with kinase, neural function, synaptotagmin and energy. These results indicate that miRNAs are substantially different between the foraging and dancing stages, and suggest that miRNAs might play important roles in regulating dancing behaviors in honey bees.

© 2012 Elsevier Ltd. All rights reserved.

## 1. Introduction

The waggle dance in honey bees (*Apis mellifera*) conveys both direction and distance information, and plays significant roles in recruiting new foragers to a food source (von Frisch, 1967; Michelsen et al., 1992). The waggle dance can also convey information about food quality (Seeley, 1995; Thom et al., 2007; Afik et al., 2008; Seefeldt and De Marco, 2008; Abbott and Dukas, 2009). In the foraging process, honey bees can integrate visual, olfactory, auditory and tactile information, and central projections of these four sensory systems involved in honey bee dance have been identified by neuronal tracer studies (Brockmann and Robinson, 2007). Recent studies have provided us with insights about the biochemical and molecular processes underlying the foraging behavior of honey bees. Octopamine is shown to initiate and maintain foraging

behavior, affect the assessment of nectar profitability (Schulz et al., 2002; Barron et al., 2007; Giray et al., 2007), and enhance appetitive learning (Hammer and Menzel, 1998). Dopamine plays a vital role in honey bees foraging, learning and motor behaviors as well (Vergoz et al., 2007; Nomura et al., 2009; Mustard et al., 2010). Since the publication of the honey bee genome, behavioral studies have been carried out at genomic, transcriptomic and proteomic levels. Foraging and dancing behaviors were affected by several genes (Johnson et al., 2002; Oldroyd and Thompson, 2006). Wang et al. (2010) analyzed the expression of the peripheral insulin receptor substrate (*IRS*) in foraging honey bees, and found that bees with reduced *IRS* expression foraged more for protein-rich foods. Nicholas (2010) discovered that distinct spatiotemporal foraging memories showed different gene expression patterns, involving nerve-related genes. Brockmann et al. (2009) found that eight peptides changed dynamically in honey bees performing different foraging behaviors.

There has been a surge of interest in recent years to determine how non-coding genes affect biological processes (Jolly and Lakhota, 2006; Mercer et al., 2009; Grillari and Grillari-Voglauer,

\* Corresponding authors. Address: College of Animal Sciences, Zhejiang University, Hangzhou 310058, China. Tel.: +86 571 88982315 (S. Su), tel.: +1 571 353 8136 (Z. Y. Huang).

E-mail addresses: [bees@msu.edu](mailto:bees@msu.edu) (Z.Y. Huang), [susongkun@zju.edu.cn](mailto:susongkun@zju.edu.cn) (S. Su).

2010). One non-coding gene Nb-1 was found to be associated with age-related division of labor in worker honey bees (Tadano et al., 2009). MicroRNAs (miRNAs) are important gene regulators that can regulate gene expressions both spatially and temporally. They can affect gene expression and the development of plants, mammals and insects by targeting mRNAs for cleavage or translational repression, and are involved in various diseases in humans (Wienholds and Plasterk, 2005; Zhang et al., 2006; Fiore et al., 2008). There have been fewer studies into honey bee miRNA: Weaver et al. (2007) verified the expression of some predicted miRNA genes experimentally in honey bees, while Chen et al. (2010) identified miRNAs in honey bees at different developmental stages utilizing SOLiD sequencing. But the roles that these miRNAs play in honey bee development or behavior have only been explored by three recent studies. Behura and Whitfield (2010) found differentially expressed miRNAs in nurses and foragers, suggesting that miRNA plays a role in regulating division of labor in honey bees. Greenberg et al. (2012) found that some specific miRNAs were not only associated with the transition from nursing to foraging, but represented a more complex relationship involving both behavioral states and colony conditions. Hori et al. (2011) studied miRNA expression in different regions of the honey bee brain, and found that some miRNAs regulate genes associated with neural regulation. However, the identified miRNAs in honey bees are limited, and their biological functions remain largely unclear.

The mechanisms underlying how miRNAs affect honey bees changing from foraging to dancing have not been investigated. There is evidence that miRNA levels can be modulated on a short term time scale (Schmitz et al., 2004; Wagner-Ecker et al., 2010). Further, miRNA is implicated in playing a role in learning and memory (Kye et al., 2011). In this study, we intended to identify miRNAs that are involved in honey bee foraging and dancing. We defined ‘dancers’ as the foraging honey bees dancing on the comb, and ‘foragers’ as the foraging honey bees who have not had a chance to exhibit dancing behavior, and are caught immediately at the entrance as they return from a foraging trip. Our hypothesis was that dancing foragers might have differentially expressed miRNAs compared to a general group of foragers, which have not had a chance to exhibit dancing behavior yet.

## 2. Materials and methods

### 2.1. Honey bee samples

Three healthy colonies of “Zhenongda-No.1” (Chen et al., 2002), a high royal jelly-yielding breed of *Apis mellifera ligustica* kept in observation hives at Zhejiang University (30.272N, 120.191E), were used in this study. Foragers and dancers ( $N = 30$  per group for both) were collected from each of three different colonies, between 11:00 and 14:00. Foragers were identified as bees returning to a colony loaded with same color and form pollen on their corbiculae, and dancers were identified as pollen foragers performing wagging dance on combs. Each forager or dancer was lowered into liquid nitrogen with a pair of forceps immediately after collection and stored at  $-80^{\circ}\text{C}$  until total RNA extraction.

### 2.2. Total RNA extraction and small RNA library construction

Ninety forager heads were pooled as the forager sample and 90 dancer heads as the dancer sample for RNA extraction. We used the method for total RNA extraction described by Liu et al. (2011). RNA extraction and small RNA library construction were performed.

For small RNA library construction, the total RNA was size fractionated on a 15% Tris–Borate–EDTA (TBE) urea polyacrylamide gel, and the 18–30 nt fraction was recovered. Small RNAs were eluted

in 0.3 M NaCl, precipitated and washed in ethanol, then dissolved in ultrapure water. The gel-purified small RNAs were ligated to the 5' RNA adapter with T4 RNA ligase, the ligation products were gel fractionated and purified before ligating to the 3' RNA adapter. The final ligation products were purified and amplified by RT-PCR. The amplification products were gel fractionated and purified by electrophoresis on a 6% polyacrylamide gel in TBE buffer, and a gel slice corresponding to the amplified library was eluted in 0.3 M NaCl. Then the purified PCR products were precipitated and washed by ethanol and resuspended in nuclease-free water. Finally, the purified PCR products were sequenced on a Solexa Sequencer (Solexa HiSeq2000, Illumina, Inc., American).

### 2.3. High-throughput sequencing and sequence analysis

After filtering out short (<18 nt), low quality and contaminant reads, the remaining reads (clean reads) were used for statistical analysis and length distribution comparison between the foragers and the dancers. The clean reads were mapped onto the *Apis mellifera* version 4.0 genome with the Short-Oligonucleotide Analysis Package (Soap, <http://www.soap.genomics.org.cn/>), and only perfectly matched tags were used for analysis.

To identify known miRNAs in each sample, we matched all the genome matched reads to non-coding sequences from NCBI Genbank database (including rRNA, tRNA, snRNA and snoRNA), Rfam database (Rfam9.0), and known miRNA hairpins of honey bee in the current release miRbase (miRBase13.0).

Because one sRNA (small RNA, RNA with 18–30 nt) may have two or more different annotations, all sRNAs were annotated in the order of rRNAetc- > known miRNA- > repeat- > exon- > intron to ensure that every sRNA has one unique annotation. The unannotated distinct small RNAs were used to identify novel candidates (sequences that have not been identified previously) with canonical hairpin structure by “Mireap” software (<http://www.sourceforge.net/projects/mireap/>).

### 2.4. Confirmation of novel miRNAs

Stem-loop RT PCR was performed to confirm the existence of novel miRNAs. Reverse transcriptase reactions were employed by using the ReverTra Ace qPCR RT Kit (TOYOBO, Japan) according to manufacturer's instructions. The miRNA-specific stem-loop primers were designed according to Chen et al. (2005). All of the stem-loop RT primers and gene-specific PCR primers are listed in Table S1. Each reaction (20  $\mu\text{l}$ ) containing 2  $\mu\text{g}$  total RNA, 4  $\mu\text{l}$   $5 \times$  RT Buffer, 1  $\mu\text{l}$  RT Enzyme Mix, 1 pM stem-loop RT primer, and nuclease-free water, was incubated in a Gradient Thermal Cycler (My Gene Series, LongGene Scientific Instruments, Co. Ltd., China) in a 96-well plate at  $37^{\circ}\text{C}$  for 15 min,  $98^{\circ}\text{C}$  for 5 min, and then  $4^{\circ}\text{C}$  for holding.

The cDNAs were diluted 1:10 to perform real-time PCR by using the THUNDERBIRD SYBR qPCR Mix (QPS-201) (TOYOBO, Japan) following manufacturer's manuals. The PCR mixture included 10  $\mu\text{l}$  THUNDERBIRD SYBR qPCR Mix, 5  $\mu\text{l}$  cDNA template, 8 pM forward primer and 8 pM reverse primer and nuclease-free water added to a final volume of 20  $\mu\text{l}$ , amplified in Line Gene K PCR system (Bioer Technology Co. Ltd., China) at  $95^{\circ}\text{C}$  for 1 min, followed by 45 cycles of 15 s at  $94^{\circ}\text{C}$  and 40 s at  $60^{\circ}\text{C}$ . The melting curve was drawn from  $60^{\circ}\text{C}$  to  $95^{\circ}\text{C}$  at a rate of  $4^{\circ}\text{C}/\text{s}$ ,  $0.5^{\circ}\text{C}$  increase per step. The melting curve for each PCR was carefully monitored to avoid non-specific amplifications. The qPCR products were detected by electrophoresis with 3% agarose gel containing Gold View and photographed by an Electrophoresis Image Analysis System (Shanghai Peiqing Science & Technology, Co. Ltd., China). Because this experiment was meant to confirm the presence of miRNAs,

we measured only one sample per miRNA and only the presence/absence of each PCR product was noted for confirmation.

### 2.5. Validation of differentially expressed miRNAs

We validated the differentially expressed miRNAs between dancers and foragers by qPCR. Thirty foragers and thirty dancers were collected from three other ZND No.1 colonies; see Section 2.1 for detailed methods. The heads of foragers and dancers from three colonies were analyzed by qPCR. The qPCR method is the same as that described in Section 2.4, except that triplicates were used for each miRNA, and quantification data were obtained. All primers used are given in Table S1. In the qPCR, U6 snoRNA was used as a control gene (Mount et al., 2007). The threshold cycle (CT) for each gene was read under the default parameters. Relative quantification of each miRNA expression was calculated by  $2^{-\Delta\Delta CT}$  (Livak and Schmittgen, 2001), and expression in foragers was used as the baseline (i.e. expression level of each miRNA in dancers was normalized to that of foragers in each colony).

### 2.6. Statistical analysis

To assess the differential expression of miRNAs of 21, 22 and 23 nt and miRNAs of other lengths between dancers and foragers, the ratio of expression levels in the two groups (dancer/forager) were calculated and *T*-tests were used to compare the average ratio in each condition to an equal-expression ratio of 1. Differences in distribution patterns in miRNA lengths between foragers and dancers were compared by contingency table analysis, and Chi Square statistic was used.

For known miRNAs expressed jointly in the two samples, the expression of miRNAs in the two samples was normalized to the same order of magnitude. We then compared their expression differences using the method described by Xin et al. (2010). We followed Xin et al. (2010), and considered the difference between expressed miRNAs to be significant only when the absolute value of fold-change ( $\log_2$  (dancer/forager)) was higher than 1, and the *P* value was less than 0.01. In the validation of differentially expressed miRNAs, *T*-tests were used as described above to determine if the differentially expressed miRNAs are significantly different.

### 2.7. Target prediction

For miRNA target gene prediction, we downloaded full-length sequences of *Apis mellifera* from the NCBI UniGene database ([ftp://ftp.ncbi.nih.gov/repository/UniGene/Apis\\_mellifera/](ftp://ftp.ncbi.nih.gov/repository/UniGene/Apis_mellifera/)), then used PITA program (Kertesz et al., 2007) to predict miRNA targets under default parameters. We selected and analyzed the target genes with  $\Delta\Delta G < -10$  kcal/mol from the initial results.

## 3. Results

### 3.1. Raw data and sequence analysis

Solexa Sequencer obtained 18,368,010 and 17,966,807 reads from foragers and dancers, respectively. After filtering out short (<18 nt), low quality and contaminant reads, we were left with 15,320,541 and 15,254,864 clean reads from foragers and dancers, respectively, which were used for further analysis.

The small RNA unique reads (the number of unique sequence types) and total reads (the number of all the reads), that were shared between foragers and dancers, and that were specific to each, are presented in Table 1.

**Table 1**  
Statistics of sRNAs from foragers and dancers.

Class	Unique	%	Total	%
Total sRNAs	2168,058	100.00	30,485,405	100.00
Shared between dancers and foragers	273,705	12.62	28,327,244	92.92
Dancer-specific	807,640	37.25	892,204	2.93
Forager-specific	1086,713	50.12	1265,957	4.15

The majority of the trimmed tags in the 10–30 nt range were concentrated in 21–23 nt (Fig. 1). The percentage of 21, 22 and 23 nt length sequences in dancers was higher than that in foragers (mean  $\pm$  SE of dancer/forager ratio =  $1.13 \pm 0.04$ , compared against 1,  $t = 5.08$ ,  $P = 0.037$ ), while the percentage of other length sequences in dancers was lower than that in foragers (mean  $\pm$  SE of dancer/forager ratio =  $0.78 \pm 0.12$ , compared against 1,  $t = -5.85$ ,  $P < 0.01$ ). The difference in distribution pattern between foragers and dancers was further supported by a significant distribution difference between the two groups when all data were analyzed together ( $\chi^2 = 322,737$ ,  $df = 27$ ,  $P < 0.0001$ ).


In both foragers and dancers, U was the predominant nucleotide at the first site of 20–24 nt length miRNAs in both samples, while A was the key nucleotide at other positions. Additionally, the A, U, C, G utilization preferences at each position were similar for foragers and dancers (Figs. S1 and S2).

Sequence analysis showed that 6694,038 reads (43.95%) in foragers and 8385,290 reads (54.97%) in dancers mapped to the genome and the total sRNAs fell into three categories: known miRNA, rRNAetc (including rRNAs, tRNAs, snRNAs and snoRNAs) and Unannotated sRNA (unannotated sRNAs).

The distributions of sRNA among the three different categories in foragers and dancers are listed in Table 2. Fifty-four known conserved miRNAs and 58 known hairpins were identified as shared between foragers and dancers. In foragers, 1270 unique reads with 3448,464 total reads can match the known hairpins, while in dancers there were 1394 unique reads with 4949,830 total reads. No miRNAs\* (complementary miRNAs) were found in our two samples (Table S2). The percentage of expression levels for each known miRNA in foragers and dancers is shown in Table S3.

### 3.2. Differential expressions of miRNAs between dancers and foragers

The expression levels of 54 miRNAs in dancers relative to foragers are presented in Table S4. Thirty-two of the 54 known miRNAs were significantly differentially expressed (with *P* value < 0.01 and an absolute value of fold-change  $\geq 1$ ). All were down-regulated in dancers. The fold-change (absolute values) of ame-miR-278,



**Fig. 1.** Length distribution of the trimmed tags in 10–30 nt range in foragers and dancers.

**Table 2**  
Distribution of small RNAs among different categories in foragers and dancers.

Category	Foragers		Dancers	
	Unique (%)	Total (%)	Unique (%)	Total (%)
Total sRNAs	1360,418 (100.00)	15,230,541 (100.00)	1081,345 (100.00)	15,254,864 (100.00)
miRNA	1270 (0.09)	3448,464 (26.64)	1394 (0.13)	4949,830 (32.45)
rRNAetc	283,211 (20.82)	6046,224 (39.70)	222,063 (20.54)	4249,302 (27.86)
Unann	1075,937 (79.09)	5735,853 (37.66)	857,888 (79.34)	6055,732 (39.70)



**Fig. 2.** Expression confirmation of novel miRNAs identified by cloning from foragers using stem-loop RT PCR.

ame-miR-210, ame-miR-282 and ame-miR-34 was over 2 (2.3-fold, 2.2-fold, 2.1-fold and 2.1-fold respectively). Ame-miR-iab-4, 133,13b, 2, 277, 317, 10, 92a, 124, 87, 14, 252, 33, 125, 929,276, 137,100, 190, 316, 1000, 927, 932, 7, 263b, 219, 981 and ame-ban-tam had a fold-change (absolute values) greater than 1 and less than 2.

### 3.3. Prediction and confirmation of novel honey bee miRNAs


Mireap predicted 86 and 104 novel miRNA candidates in foragers and dancers, respectively. There were 13 and 14 novel miRNAs with the total reads >1000 (TPM) in foragers and dancers respectively. Among these, two miRNA candidates were specific for foragers and three were specific for dancers. Eleven common and five specific novel miRNA candidates were chosen to perform Stem-loop RT PCR for confirmation and quantification. All the 16 novel miRNA sequences are shown in Table S5. PCR results showed that all of them yielded PCR products (Fig. 2), which indicated that these novel miRNAs were expressed in both foragers and dancers.

### 3.4. Validation of differential miRNA expression by qPCR

Solexa sequencing results showed that most of miRNAs were down-regulated in dancers. We selected the top 4 miRNAs with significant differences in expression (the absolute value of fold-change was larger than 2) for validation by qPCR. Two known miRNAs, ame-miR-278 and ame-miR-282, were significantly down regulated in dancers compared to foragers ( $t = -8.89$ ,  $df = 2$ ,  $P < 0.05$ , and  $t = -15.66$ ,  $df = 2$ ,  $P < 0.01$ , respectively, the average ratio of dancer expressions/forager expressions compared to an equal-expression ratio of 1) (Fig. 3).

### 3.5. Target prediction

We used the PITA program to predict the target genes of ame-miR-278 and ame-miR-282, each with thousands of targets. We selected potential targets taking both  $ddG < -10$  kcal/mol and the related functions into consideration. High-probability targets of ame-miR-278 and ame-miR-282 are shown in Table 3. Targets of ame-miR-278 included *Apis mellifera* protein kinase C



**Fig. 3.** Validation of specific down-regulated miRNA genes in dancers compared to foragers in three new colonies.

(NM\_001134948, Pkc), *Apis mellifera* NADH dehydrogenase (ubiquinone) Fe-S protein 1, 75 kDa (NADH-coenzyme Q reductase) (NM\_001172404, Ndufs1), *Apis mellifera* synaptotagmin 20 (NM\_001142561, Syt20), *Apis mellifera* mitochondrial ribosomal protein L2 (XM\_625016, mRpl2) and so on. The four important targets of ame-miR-282 were *Apis mellifera* protein kinase C (NM\_001134948, Pkc), *Apis mellifera* synaptotagmin 20 (NM\_001142561, Syt20), *Apis mellifera* Tie-like receptor tyrosine kinase (XM\_393849, Tie) and *Apis mellifera* neurogenic locus protein delta (XM\_393831, Dl). Most of the target genes were associated with kinase, neural function, synaptotagmin and energy.

## 4. Discussion

In this study, Solexa sequencing was used to compare miRNA differences in foragers and dancers. The sequencing results showed that the types of shared sRNA (common small RNA unique reads) in foragers and dancers were 12.62% of the total unique reads, while their expression accounted for 92.92% of the total reads. This suggests that a small number of miRNAs play fundamental roles in honey bees for maintaining basic life processes common to both foragers and dancers.

The specific unique reads in dancers were 37.25% of the total unique reads, whereas their expression was only 2.93% of the total reads (Table 1). The same trend was found in foragers. It is possible that only a few miRNAs are needed to have high expression levels. Many studies have pointed that one miRNAs could target hundreds of mRNAs via imprecise base pairing (Bartel and Chen, 2004; Grun et al., 2005; Bartel, 2009), so that a few miRNAs could have significant effects on post-transcriptional regulation. In addition, some miRNAs can work together through synergism (Bartel, 2009).

Jagadeeswaran et al. (2010) found that the length distribution of total reads is bimodal, with one group peaking around 20–22 nt (miRNA) and another around 26–29 nt (piRNA). Our study only found one peak with lengths of 21–23 nt in honey bees. The first nucleotide bias and the nucleotide bias at each position of both known miRNAs and novel miRNAs were similar between for-



**Table 3**  
High-probability targets of ame-miR-278 and ame-miR-282.

miRNAs	Predicted targets
ame-miR-278	Pkc, Imd, Ndufs1, Dscam, Evg, Syt20, LOC724697, Pfrx, LOC410662, ec, mRpL2, CUBN, Blimp-1, LOC408646, Atg2
ame-miR-282	Pkc, Syt20, LOC551139, LOC552057, LOC724697, Tie, DI, LOC552646, LOC408820, LOC409821, LOC725936, LOC724430, LOC412777

agers and dancers, indicating the stability of miRNA sequence structures, a factor which is important for functional maintenance (Khvorova et al., 2003; Girard et al., 2006). The first and the whole nucleotide bias of novel miRNAs were different from that of known miRNAs (Figs. S1 and S2). This indicates that these novel miRNAs have different structures from known miRNAs, and may have different functions. Total sRNAs mapped to genome could be placed in three categories, namely known miRNA, rRNA etc and the Unannotated sRNA. Table 2 shows that in foragers and dancers, the unique reads of unannotated sRNAs and rRNAs were much more numerous than those of miRNAs, while the total reads of unannotated sRNAs, rRNAs and miRNAs were almost equal. These data suggest that although miRNA unique reads only accounted for a small percentage of sRNA unique reads, they compensated for this through higher expression. Furthermore, there might be important novel miRNAs among the un-annotated sRNAs.

The following miRNAs were the 5 miRNAs with the highest expression levels (total reads) both in foragers and dancers: ame-miR-1 (69.64%), ame-miR-276 (7.87%), ame-miR-275 (4.20%), ame-miR-184 (3.14%) and ame-miR-996 (2.58%) (Table S3), and these may be associated with fundamental vital functions. Some studies found that miR-1 is involved in muscle development (Nguyen and Frasch, 2006) and the Notch pathway, which is vital for the development of the peripheral nervous system in *Drosophila* (Kwon et al., 2005; Wheeler et al., 2008). Two miRNAs, mir-1 and mir-133 were found to be highly conserved and co-located in adjacent gene locations in insects and vertebrates (Rao et al., 2006; Yu et al., 2008), which is consistent with our data showing that both were highly expressed in foragers and dancers. Ame-miR-276 was preferentially expressed in the olfactory lobes and the class I and II small-type Kenyon cells (cells involved in learning and memory in honey bees, Szyszka et al., 2008) in the honey bee brain, and ame-miR-276 and -1000 could target the neurally related genes (Hori et al., 2011). Furthermore, it has been suggested that miR-184 is involved in the central nervous system in *Drosophila* (Li et al., 2011). These studies, along with our results, indicate that the highly expressed miRNAs, ame-miR-1, ame-miR-276 and ame-miR-184 may be involved in neural function and perhaps in foraging and dancing behaviors.

Solexa sequencing data showed that the majority of miRNAs were down-regulated in dancers, and we further confirmed that the expression of ame-miR-278 and ame-miR-282 were lower in dancers than in foragers, suggesting that the target genes of these miRNAs are activated in dancers. Two studies revealed that in *Drosophila* and the silkworm, lower miR-278 expression is associated with a higher metabolic rate (Teleman et al., 2006; Jagadeeswaran et al., 2010). The lower expression of ame-miR-278 found in dancers, therefore, suggests that dancers might have a higher metabolic rate. In target prediction, most target genes of ame-miR-278 and ame-miR-282 were associated with kinase (such as Pkc and Pfrx), neural function (such as Tie and DI), synaptotagmin (such as Syt20, CUBN and LOC552057) and energy (such as Ndufs1 and mRpL2), suggesting that these genes play important functions in bee dancing behaviors. The functions of ame-miR-278 and ame-miR-282 need experimental verification through further behavioral studies. Some non-coding genes, such as *Ks-1* and *Nb-1*, were shown to have a possible role in modulating honey bee behaviors (Sawata et al., 2002; Tadano et al., 2009). MiR-8 has been found

to regulate neuromuscular junction formation (Loya et al., 2009) and participate in the Wingless/Wnt pathway (Kennell et al., 2008), which is associated with brain development in *Drosophila*. Recently, a hypothetical miR-8 binding site was found in the honey bee Wingless gene (Hori et al., 2011). These studies strongly suggest that more miRNAs and more non-coding genes associated with foraging and dancing behaviors will be found in the future.

In our study, we tried to find differentially expressed miRNAs between foragers and dancers; naturally, these are groups of worker bees at different stages of the foraging process. In order to overcome individual differences (including colony, age, genetics and so on), we collected foragers and dancers one by one in turn from the same colony at the same time and pooled samples from three colonies together. Thus, we established two representative sRNA pools of heads from foragers and dancers, using a powerful next-generation sequencing method. These sRNA pools contained all sRNAs involving in foraging behavior and dancing behavior completely. Comparing the two sRNA pools, we found some differences in miRNAs in the two behavioral groups, and confirmed the differential expressions of several miRNAs between the two behavioral groups, using bees from three new colonies. The results showed that miRNAs play important roles in the process of switching from foraging to dancing. More studies are needed to discover the molecular mechanism underlying the dancing behavior in the future.

### Acknowledgements

The research was supported by earmarked funds for Modern Agro-industry Technology Research System (No. CARS-45-KXJ3) and Nature and Science Foundation Commission of Zhejiang Province (R3080306) to SKS. The deep sequencing and bio-informatics analysis work were carried out in Beijing Genome Institute (<http://www.genomics.cn/en/index.php>). We thank Dr. Aung Si and two anonymous referees for helpful suggestions that improved the manuscript, Shuangjin Dai for his beekeeping, Yunjuan Hu and Qin Chen for their help in research.

### Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.jinsphys.2012.08.008>.

### References

- Abbott, K.R., Dukas, R., 2009. Honeybees consider flower danger in their waggle dance. *Animal Behaviour* 78, 633–635.
- Afik, O., Dag, A., Shafir, S., 2008. Honeybee, *Apis mellifera*, round dance is influenced by trace components of floral nectar. *Animal Behaviour* 75, 371–377.
- Barron, A.B., Maleszka, R., Vander Meer, R.K., Robinson, G.E., 2007. Octopamine modulates honey bee dance behavior. *Proceedings of the National Academy of Sciences* 104, 1703–1707.
- Bartel, D.P., 2009. MicroRNAs: target recognition and regulatory functions. *Cell* 136, 215–233.
- Bartel, D.P., Chen, C.Z., 2004. Micromanagers of gene expression: the potentially widespread influence of metazoan microRNAs. *Nature Reviews Genetics* 5, 396–401.
- Behura, S.K., Whitfield, C.W., 2010. Correlated expression patterns of microRNA genes with age-dependent behavioural changes in honeybee. *Insect Molecular Biology* 19, 431–439.

- Brockmann, A., Robinson, G.E., 2007. Central projections of sensory systems involved in honey bee dance language communication. *Brain, Behavior and Evolution* 70, 125–136.
- Brockmann, A., Annangudi, S.P., Richmond, T.A., 2009. Quantitative peptidomics reveal brain peptide signatures of behavior. *Proceedings of the National Academy of Sciences* 106, 2383–2388.
- Chen, S.L., Su, S.K., Lin, X.Z., 2002. An introduction to high-yielding royal jelly production methods in China. *Bee World* 83, 69–77.
- Chen, C., Ridzon, D.A., Broomer, A.J., Zhou, Z., Lee, D.H., Nguyen, J.T., Barbisin, M., Xu, N.L., Mahuvakar, V.R., Andersen, M.R., Lao, K.Q., Livak, K.J., Guegle, K.J., 2005. Real-time quantification of microRNAs by stem-loop RT-PCR. *Nucleic Acids Research* 33, e179.
- Chen, X., Yu, X., Cai, Y., Zheng, H., Yu, D., Liu, G., Zhou, Q., Hu, S., Hu, F., 2010. Next-generation small RNA sequencing for microRNAs profiling in the honey bee *Apis mellifera*. *Insect Molecular Biology* 19, 799–805.
- Fiore, R., Siegel, G., Schratt, G., 2008. MicroRNA function in neuronal development, plasticity and disease. *Biochimica et Biophysica Acta* 1779, 471–478.
- Girard, A., Sachidanandam, R., Hannon, G.J., Carmell, M.A., 2006. A germline-specific class of small RNAs binds mammalian Piwi proteins. *Nature* 442, 199–202.
- Giray, T., Galindo-Cardona, A., Oskay, D., 2007. Octopamine influences honey bee foraging preference. *Journal of Insect Physiology* 53, 691–698.
- Greenberg, J.K., Xia, J., Zhou, X., Thatcher, S.R., Gu, X., Ament, S.A., Newman, T.C., Green, P.J., Zhang, W., Robinson, G.E., Ben-Shahar, Y., 2012. Behavioral plasticity in honey bees is associated with differences in brain microRNA transcriptome. *Genes, Brain and Behavior*. <http://dx.doi.org/10.1111/j.1601-183X.2012.00782.x>.
- Grillari, J., Grillari-Voglauer, R., 2010. Novel modulators of senescence, aging, and longevity: small non-coding RNAs enter the stage. *Experimental Gerontology* 45, 302–311.
- Grun, D., Wang, Y.L., Langenberger, D., Gunsalus, K.C., Rajewsky, N., 2005. MicroRNA target predictions across seven *Drosophila* species and comparison to mammalian targets. *PLoS Computational Biology* 1, e13.
- Hammer, M., Menzel, R., 1998. Multiple sites of associative odor learning as revealed by local brain microinjections of octopamine in honeybees. *Learning and Memory* 5, 146–156.
- Hori, S., Kaneko, K., Saito, T.H., Takeuchi, H., Kubo, T., 2011. Expression of two microRNAs, ame-mir-276 and -1000, in the adult honeybee (*Apis mellifera*) brain. *Apidologie* 42, 89–102.
- Jagadeeswaran, G., Zheng, Y., Sumathipala, N., Jiang, H., Arrese, E.L., Soulages, J.L., Zhang, W., Sunkar, R., 2010. Deep sequencing of small RNA libraries reveals dynamic regulation of conserved and novel microRNAs and microRNA-stars during silkworm development. *BMC Genomics* 11, 52.
- Johnson, R.N., Oldroyd, B.P., Barron, A.B., Crozier, R.H., 2002. Genetic control of the honey bee (*Apis mellifera*) dance language: segregating dance forms in a backcrossed colony. *The Journal of Heredity* 93, 170–173.
- Jolly, C., Lakhotia, S.C., 2006. Human sat III and *Drosophila* hsr $\alpha$  transcripts: a common paradigm for regulation of nuclear RNA processing in stressed cells. *Nucleic Acids Research* 34, 5508–5514.
- Kennell, J.A., Gerin, I., MacDougald, O.A., Cadigan, K.M., 2008. The microRNA miR-8 is a conserved negative regulator of Wnt signaling. *Proceedings of the National Academy of Sciences* 105, 15417–15422.
- Kertesz, M., Iovino, N., Unnerstall, U., Galau, U., Segal, E., 2007. The role of site accessibility in microRNA target recognition. *Nature Genetics* 39, 1278–1284.
- Khvorova, A., Reynolds, A., Jayasena, S.D., 2003. Functional siRNAs and miRNAs exhibit strand bias. *Cell* 115, 209–216.
- Kwon, C., Han, Z., Olson, E.N., Srivastava, D., 2005. MicroRNA1 influences cardiac differentiation in *Drosophila* and regulates Notch signaling. *Proceedings of the National Academy of Sciences* 102, 18986–18991.
- Kye, M.J., Neveu, P., Lee, Y.S., Zhou, M., Steen, J.A., Sahin, M., Kosik, K.S., Silva, A.J., 2011. NMDA mediated contextual conditioning changes miRNA expression. *PLoS One* 6, e24682. <http://dx.doi.org/10.1371/journal.pone.0024682>.
- Li, P., Peng, J., Hu, J., Xu, Z., Xie, W., Yuan, L., 2011. Localized expression pattern of miR-184 in *Drosophila*. *Molecular Biology Reports* 38, 355–358.
- Liu, F., Li, W.F., Li, Z.G., Zhang, S.W., Chen, S.L., Su, S.K., 2011. High-abundance mRNAs in *Apis mellifera*: comparison between nurses and foragers. *Journal of Insect Physiology* 57, 274–279.
- Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>- $\Delta\Delta$ CT</sup> Method. *Methods* 25, 402–408.
- Loya, C.M., Lu, C.S., Vactor, D.V., Fulga, T.A., 2009. Transgenic microRNA inhibition with spatiotemporal specificity in intact organisms. *Nature Methods* 6, 897–903.
- Mercer, T.R., Dinger, M.E., Mattick, J.S., 2009. Long non-coding RNAs: insights into functions. *Nature Reviews Genetics* 10, 155–159.
- Michelsen, A., Andersen, B.B., Storm, J., Kirchner, W.H., Lindauer, M., 1992. How honeybees perceive communication dances, studied by means of a mechanical model. *Behavioral Ecology and Sociobiology* 30, 143–150.
- Mount, S.M., Gotea, V., Lin, C., Hernandez, K., Makalowski, W., 2007. Spliceosomal small nuclear RNA genes in 11 insect genomes. *RNA* 13, 5–14.
- Mustard, J.A., Pham, P.M., Smith, B.H., 2010. Modulation of motor behavior by dopamine and the D1-like dopamine receptor AmDOP2 in the honey bee. *Journal of Insect Physiology* 56, 422–430.
- Nguyen, H.T., Frasch, M., 2006. MicroRNAs in muscle differentiation: lessons from *Drosophila* and beyond. *Current Opinion in Genetics and Development* 16, 533–539.
- Nicholas, L.N., 2010. Spatiotemporal Memories in Honey Bees: Microarray Analysis of Time-Trained Foragers. M.S. Thesis, Urbana, Illinois, University of Illinois, p. 69. [http://www.ideals.illinois.edu/bitstream/handle/2142/16789/2\\_Naeger%20Nicholas.pdf?sequence=4](http://www.ideals.illinois.edu/bitstream/handle/2142/16789/2_Naeger%20Nicholas.pdf?sequence=4) (visited 13.10.2011).
- Nomura, S., Takahashi, J., Sasaki, T., Yoshida, T., Sasaki, M., 2009. Expression of the dopamine transporter in the brain of the honeybee, *Apis mellifera* L. (Hymenoptera: Apidae). *Applied Entomology and Zoology* 44, 403–411.
- Oldroyd, B.P., Thompson, G.J., 2006. Behavioural genetics of the honey bee *Apis mellifera*. *Advances in Insect Physiology* 33, 1–49.
- Rao, P.K., Kumar, R.M., Farkhondeh, M., Baskerville, S., Lodish, H.F., 2006. Myogenic factors that regulate expression of muscle-specific microRNAs. *Proceedings of the National Academy of Sciences* 103, 8721–8726.
- Sawata, M., Yoshino, D., Takeuchi, H., Kamikouchi, A., Ohashi, K., Kubo, T., 2002. Identification and punctate nuclear localization of a novel noncoding RNA, *Ks-1*, from the honeybee brain. *RNA* 8, 772–785.
- Schmitz, S., Pfaffl, M.W., Meyer, H.H.D., Bruckmaier, R.M., 2004. Short-term changes of mRNA expression of various inflammatory factors and milk proteins in mammary tissue during LPS-induced mastitis. *Domestic Animal Endocrinology* 26, 111–126.
- Schulz, D.J., Barron, A.B., Robinson, G.E., 2002. A role for octopamine in honey bee division of labor. *Brain, Behavior and Evolution* 160, 350–359.
- Seefeldt, S., De Marco, R.J., 2008. The response of the honeybee dance to uncertain rewards. *Journal of Experimental Biology* 211, 3392–3400.
- Seeley, T.D., 1995. *The Wisdom of the Hive: The Social Physiology of Honey Bee Colonies*. Harvard University Press, Cambridge, p. 295.
- Szyska, P., Galkin, A., Menzel, R., 2008. Associative and non-associative plasticity in Kenyon cells of the honeybee mushroom body. *Frontiers in Systems Neuroscience* 2, 3. <http://dx.doi.org/10.3389/neuro.06.003.2008>.
- Tadano, H., Yamazaki, Y., Takeuchi, H., Kubo, T., 2009. Age- and division-of-labour-dependent differential expression of a novel non-coding RNA, *Nb-1*, in the brain of worker honeybees, *Apis mellifera* L. *Insect Molecular Biology* 8, 715–726.
- Teleman, A.A., Maitra, S., Cohen, S.M., 2006. *Drosophila* lacking microRNA miR-278 are defective in energy homeostasis. *Genes and Development* 20, 417–422.
- Thom, C., Gilley, D.C., Hooper, J., Esch, H.E., 2007. The scent of the waggle dance. *PLoS Biology* 5, 1862–1867.
- Vergoz, V., Roussel, E., Sandoz, J.C., Giurfa, M., 2007. Aversive learning in honeybees revealed by the olfactory conditioning of the sting extension reflex. *PLoS One* 2, e288.
- von Frisch, K., 1967. *The Dance Language and Orientation of Bees*. Harvard University Press, Cambridge (Massachusetts), p. 161.
- Wagner-Ecker, M., Schwager, C., Wirkner, U., Abdollahi, A., Huber, P.E., 2010. MicroRNA expression after ionizing radiation in human endothelial cells. *Radiation Oncology* 5, 25. <http://dx.doi.org/10.1186/1748-717X-5-25>.
- Wang, Y., Mutti, N.S., Ihle, K.E., Siegel, A., Dolezal, A.G., Kaftanoglu, O., Amdam, G.V., 2010. Down-regulation of honey bee *IRS* gene biases behavior toward food rich in protein. *PLoS Genetics* 6, e1000896.
- Weaver, D.B., Anzola, J.M., Evans, J.D., Reid, J.G., Reese, J.T., Childs, K.L., Zdobnov, E.M., Samanta, M.P., Miller, J., Elsik, C.G., 2007. Computational and transcriptional evidence for microRNAs in the honey bee genome. *Genome Biology* 8, R97. <http://dx.doi.org/10.1186/gb-2007-8-6-r97>.
- Wheeler, S.R., Stagg, S.B., Crews, S.T., 2008. Multiple Notch signaling events control *Drosophila* CNS midline neurogenesis, gliogenesis and neuronal identity. *Development* 135, 3071–3079.
- Wienholds, E., Plasterk, R.H.A., 2005. MicroRNA function in animal development. *FEBS Letters* 579, 5911–5922.
- Xin, M., Wang, Y., Yao, Y., Xie, C., Peng, H., Ni, Z., Sun, Q., 2010. Diverse set of microRNAs are responsive to powdery mildew infection and heat stress in wheat (*Triticum aestivum* L.). *BMC Plant Biology* 10, 123. <http://dx.doi.org/10.1186/1471-2229-10-123>.
- Yu, X., Zhou, Q., Li, S.C., Luo, Q., Cai, Y., Lin, W., Chen, H., Yang, Y., Hu, S., Yu, J., 2008. The silkworm (*Bombyx mori*) microRNAs and their expressions in multiple developmental stages. *PLoS One* 3, e2997.
- Zhang, B., Pan, X., Cobb, G.P., Anderson, T.A., 2006. Plant microRNA: a small regulatory molecule with big impact. *Developmental Biology* 289, 3–16.